


AD _____

GRANT NO:
DAMD17-94-J-4473



TITLE:

Estrogen Responsive Breast Cancer Growth Regulation

PRINCIPAL INVESTIGATOR:

David A. Sirbasku, Ph.D.

CONTRACTING ORGANIZATION:

The University of Texas Health Science Center at Houston, Texas 77225

REPORT DATE:

October 5, 1995

TYPE OF REPORT:

Annual Report

19951220 048

PREPARED FOR: U.S. Army Medical Research and Materiel
Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 3

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 5, 1995		3. REPORT TYPE AND DATES COVERED Annual 15 Sep 94 - 14 Sep 95
4. TITLE AND SUBTITLE Estrogen Responsive Breast Cancer Growth Regulation			5. FUNDING NUMBERS DAMD17-94-J-4473	
6. AUTHOR(S) David A. Sirbasku, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Texas Health Sciences Center at Houston Houston, Texas 77225			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Not Applicable				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) In this U.S. Army sponsored research program, we proposed to identify and characterize the serum factor(s) which regulated estrogen-responsive breast cancer cell growth and to conduct another study to ask if loss of serum factor(s) control was related to hormone autonomy. During the first year, we have been successful beyond our expectations. We have demonstrated that two well known plasma glycoproteins, sex hormone-binding globulin (SHBG) and corticosteroid binding globulin (CBG) are negative regulators of steroid hormone responsive growth and that estrogens and progesterone reverse this inhibition. Furthermore, steroid hormone autonomous cells appear not to be regulated by either SHBG or CBG. Our protein chemistry data show that the SHBG we have isolated is not the common (Type I) form most often studied, but a variant that has only 30 to 40% amino acid sequence homology. The results to date suggest that breast cancer growth may be regulated by a new form of SHBG which we have designated Type II. The search for human Type II SHBG will be a major topic of this research for the next year.				
14. SUBJECT TERMS Estrogens; Progestins; Cell Growth; Inhibitors; breast cancer Sex Hormone-Binding Globulin; Corticosteroid-Binding Globulin			15. NUMBER OF PAGES 20	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to *stay within the lines* to meet optical scanning requirements.

Block 1. Agency Use Only (Leave blank).

Block 2. Report Date. Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.

Block 3. Type of Report and Dates Covered. State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 - 30 Jun 88).

Block 4. Title and Subtitle. A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.

Block 5. Funding Numbers. To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

C - Contract	PR - Project
G - Grant	TA - Task
PE - Program Element	WU - Work Unit Accession No.

Block 6. Author(s). Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).

Block 7. Performing Organization Name(s) and Address(es). Self-explanatory.

Block 8. Performing Organization Report Number. Enter the unique alphanumeric report number(s) assigned by the organization performing the report.

Block 9. Sponsoring/Monitoring Agency Name(s) and Address(es). Self-explanatory.

Block 10. Sponsoring/Monitoring Agency Report Number. (If known)

Block 11. Supplementary Notes. Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of...; To be published in.... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

Block 12a. Distribution/Availability Statement. Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

DOD - See DoDD 5230.24, "Distribution Statements on Technical Documents."

DOE - See authorities.

NASA - See Handbook NHB 2200.2.

NTIS - Leave blank.

Block 12b. Distribution Code.

DOD - Leave blank.

DOE - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.

NASA - Leave blank.

NTIS - Leave blank.

Block 13. Abstract. Include a brief (*Maximum 200 words*) factual summary of the most significant information contained in the report.

Block 14. Subject Terms. Keywords or phrases identifying major subjects in the report.

Block 15. Number of Pages. Enter the total number of pages.

Block 16. Price Code. Enter appropriate price code (*NTIS only*).

Blocks 17. - 19. Security Classifications. Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.

Block 20. Limitation of Abstract. This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

N/A Where copyrighted material is quoted, permission has been obtained to use such material.

N/A Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

N/A Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

DAS DAS In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

N/A For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

DAS DAS In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

DAS DAS In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

David A. Subash 10/05/95
PI - Signature Date

PI: David A. Sirbasku, Ph.D.

I. TABLE OF CONTENTS

CONTENTS OF THE ANNUAL REPORT	PAGE NUMBER
Front cover	1
SF 298 Report Documentation Page	2
Foreword	3
Table of Contents	4
Introduction	5 - 10
Body	10 - 15
Conclusions	15 - 16
References	16 - 20
Appendix	None

PI: David A. Sirbasku, Ph.D.

II. INTRODUCTION

A. Nature of the Problem

During the first 12 months of our US Army Breast Cancer Program project, we have successfully addressed one of the major unresolved issues in breast cancer endocrine physiology. In our original proposal, we planned first (TASK 1) to identify the serum factor(s) which regulated estrogen-dependent breast cancer cell growth. We are pleased to report that two exceptionally significant serum factors have been identified. They both are well known human plasma glycoproteins. Both are related to the action of sex steroids or progestins. During the first year of this project, we purified these serum factors from horse and rat serum and characterized them by several different methods including partial amino acid sequencing.

B. Background

This project is based on recent discoveries in our laboratory which have resolved a longstanding problem in breast cancer endocrine physiology. In 1896, Beatson reported that oophorectomy had palliative effects on breast cancer (1). That report first implicated estrogens and progestins in breast cancer growth. Since that time, researchers have identified specific nuclear receptors for these steroid hormones (2-3) and characterized them by cDNA cloning and many other methods (4). This advance led to the assumption that the interaction of estrogens and progestins with their specific nuclear receptors was sufficient to explain target tissue growth (2-3). However, as work progressed, this point of view became less defensible.

Many investigators came to the conclusion that other regulatory molecules also were required. For example, our laboratory proposed that plasma contained an estrogen regulated mediating "estromedin" which controlled target cell proliferation (5). Several putative estromedins were purified from tissues and identified as insulin-like growth factors I and II (IGF-I and IGF-II) (6,7), transforming growth factor α (TGF α) (8), and fibroblast growth factors (9). None proved to be related to estrogen dependent breast cancer cell growth (10). Another laboratory proposed that plasma contained an inhibitor (estroclyone) which negatively controlled target tissue growth (11). It has been proposed that estroclyone inhibits growth by binding to target cells, and that this restraint is reversed when steroid hormones associate with the inhibitor to release it from cells. Estroclyones have proven to be unstable during isolation and because of this have not been characterized at the molecular level (12). Another group concluded that serum albumin was "estroclyone" (13). This hypothesis was not confirmed (10). Continuing with the theme of serum factor regulation, other investigators did not offer specific names for the agents involved, but did conclude that serum contained a critically important regulator(s) which had not yet been identified (14-17). In support of this viewpoint, it was known that estrogen mitogenic effects *in vitro* were difficult to demonstrate (17-20), especially in completely serum-free hormonally defined medium (10,21,22). When estrogen mitogenic effects were found *in vitro*, they were most pronounced in serum supplemented cultures (11,17,23-26). Recently, we found that the ferric iron in culture medium was deleterious to hormone-responsive growth (27). Use of "low-iron" medium now

PI: David A. Sirbasku, Ph.D.

permits identification of very large estrogen mitogenic effects in serum-supplemented culture (27).

The work of Sporn and Todaro (28) opened new possibilities with regard to growth factors and cancer. They proposed that growth factors might act as autocrine, paracrine or endocrine regulators. Many groups joined in these studies. For example, one consortium of investigators proposed that estrogens induced TGF α , IGF-I and/or IGF-II, and that these acted as autocrine, paracrine or intracrine agents to promote breast cancer growth (29-35). Despite the attractiveness of these hypotheses, such mechanisms have yet to be established conclusively. Data from the original laboratories and others (36-39), and from our group (10), have raised serious questions concerning participation of these growth factors in estrogen-dependent growth.

Our recent work supported by US Army grant DAMD17-94-J-4473 has now resolved this issue. We have discovered that two well known and thoroughly characterized plasma glycoproteins, SHBG and CBG, are negative regulators (*i.e.* inhibitors) of estrogen receptor positive (ER⁺) and progesterone receptor positive (PR⁺) breast cancer cell growth, respectively, and that autonomous (ER⁻ and PR⁻) breast cancer cells are not inhibited by these glycoproteins. The experimental results describing these studies are part of a series of nine manuscripts in preparation (see the listing below)

SHBG, which is a M_r 94,000 homodimer (40,41), has been identified as the major plasma carrier of sex hormones (42-45). CBG is a M_r 58,000 monomer which serves that function for progestins and corticosteroids (42). These proteins appeared to have no function beyond regulating the relative levels of "free/active" versus "bound/inactive" hormones in plasma (42,46). Evidence suggesting direct cellular functions became apparent only after the identification of SHBG receptors on sex hormone target tissues (47-57) and CBG receptors on human breast cancer and other target cells (42,58,59,61). Nevertheless, the function of these sites has remained unexplained. Studies have asked if these receptors might be docking sites for delivery of steroid hormones to the interior of cells. However, further work suggests this is unlikely (42,49). More positively, the receptors appear to possess a transmembrane signaling function which implies a regulatory role. When hormone free SHBG or CBG are bound to their receptors, the association of an appropriate steroid with the bound ligand induces a transient increase in cyclic AMP production (42,55,59,60).

Most of what is known about SHBG and CBG receptors has come from measuring the binding of ¹²⁵I-labeled SHBG or ¹²⁵I-CBG to whole cells, membrane fractions of cells or detergent solubilized membranes. The solubilized SHBG receptor from prostate cells showed an ¹²⁵I-SHBG affinity of K_a ~ 3 x 10¹⁰ M⁻¹ (54,55). Studies with ¹²⁵I-CBG and human breast cancer MCF-7 cells gave K_a of ~ 1.4 x 10⁹ M⁻¹ (42,59). Studies have estimated the M_r of the SHBG receptor at ~ 170,000 (52,54). Another study showed that the amino acid residues 48-57 of SHBG contained a receptor recognition sequence (56). Beyond this information however, not even partial amino acid sequences are available for these receptors.

To place our study in perspective, it is vital to recall that the loss of negative regulation already has been implicated in a striking increase in susceptibility to breast cancer. For example, the autosomal dominant BRCA1 gene appears to code for a tumor suppressor protein (62-64). Mutations in BRCA1 confer a risk of 85% to breast and ovarian cancer. This autosomal dominant trait is carried by about 5% of the women in the United States. Also, the loss of the function (*i.e.* mutation) of the p53 tumor suppressor protein has important implications for germline and sporadic (*i.e.* non-familial) breast cancer susceptibility (65-68).

PI: David A. Sirbasku, Ph.D.

C. Purpose of the Present Work

Shown below is our original Statement of Work. The purpose of the next year will be to continue with the primary goal of our original proposed research, namely, to structurally define the serum factor(s) which regulates steroid hormone dependent breast cancer cell growth. While these studies already have yielded remarkable new insights, they have also expanded Task 1 to include additional purification work and more cDNA cloning and DNA sequencing. Before our discoveries this past year, SHBG and CBG had never been implicated in breast cancer growth. Indeed, to make matters even more interesting, the form of SHBG we have isolated was completely unexpected. *These discoveries bear directly on our original theme of the negative regulation of breast cell growth.*

G. STATEMENT OF WORK

TITLE: ESTROGEN RESPONSIVE BREAST CANCER GROWTH REGULATION

TASK 1. Purification of Estrogen Mediating Serum Factor (EMSF): Months 12-30

- (a) Purification will be done by application of conventional, FPLC and HPLC methods using horse and, if indicated, human serum.
- (b) N^o and internal amino acid microsequencing will be done.
- (c) Computer registered sequences will be used to identify EMSF.
- (d) Rabbit polyclonal antisera will be raised against ovalbumin conjugates of EMSF peptides synthesized by solid phase methods.
- (e) Immunoabsorption will be used to confirm the relationship between amino acid sequence and activity.
- (f) The tissue of origin of EMSF will be sought in mouse organ extracts by Northern blotting for m-RNA.
- (g) The effects of estrogen status on EMSF concentration in mouse blood will be assessed by ELISA or RIA.
- (h) cDNA libraries will be screened for EMSF containing plasmids.
- (i) If the complete sequence of EMSF is not known, we will determine it by dsDNA dideoxy chain termination sequencing.

TASK 2. Characterization of the EMSF Receptor: Months 12-48

- (a) ¹²⁵I-EMSF will be made by the chloramine T reaction.
- (b) Specific ¹²⁵I-EMSF binding will be investigated with estrogen sensitive breast cancer cells. The experimental parameters will be time, concentration, temperature, and effects of estrogens in the culture medium. Scatchard analysis is expected to give K_d values and numbers of sites per cell.
- (c) Chemical cross-linking with ¹²⁵I-EMSF followed by SDS-PAGE and autoradiography will be used to identify the M_r of the specific ligand binding receptor.
- (d) Internalization will be studied to determine if EMSF undergoes receptor mediated endocytosis.
- (e) The same studies outlined in (a) will be done to determine if estrogen insensitive breast cancer cells lack EMSF specific binding.
- (f) MAb's will be raised which block ¹²⁵I-EMSF binding to receptors.
- (g) The MAb's will be used to screen cDNA libraries from estrogen sensitive cells to obtain plasmids for dideoxy dsDNA sequencing of the complete receptor.
- (h) Transfection with full length receptor cDNA will be evaluated to determine if estrogen insensitive cells can be reverted to negative control by EMSF.

TASK 3. Physiological Significance in vitro and in vivo: Months 30-48

- (a) The MAb's raised will be evaluated as growth agonists and antagonists in serum-free culture with estrogen sensitive breast cancer cells.
- (b) The effects of antagonistic MAb's will be investigated on estrogen sensitive and insensitive breast cancer cell tumor formation athymic nude mice.

PI: David A. Sirbasku, Ph.D.

Human SHBG has been thoroughly characterized. It is purified from pregnant human female plasma or from female plasma by androgen (testosterone) affinity chromatography. This kind of SHBG is known in many species. We have tentatively designated it *Type I human SHBG* based on its amino acid sequence and its high affinity for sex steroid hormones. We have isolated a new form of SHBG from horse serum and rat serum. This new type does not share high sequence homology with the testosterone affinity purified SHBGs from human and rabbit. In fact, the homology is at best 30 to 40%. Also, horse and rat SHBG bind androgens and estrogens with a ten-fold lower affinity than Type I human SHBG. The marked contrasts in primary structure apparently cause differences in antigenicity and steroid hormone binding. These variant characteristics may explain why Type II SHBG has not been identified before.

Based on these new discoveries, the statement of work for the next year is described next:

CONTINUATION TASKS FOR THE UPCOMING NEXT YEAR (1995-1996):
(1) We will continue the identification of Type II SHBG by performing cDNA cloning and sequencing of the horse and rat proteins (an expanded form of original tasks 1h and 1i).
(2) We will use both biochemical isolation and molecular cloning methods to seek human Type II SHBG (part of original task 1a).
(3) We will develop immunological methods of measuring Type II SHBG in rat plasma and human serum samples (part of original tasks 1d and 1e).
(4) We will determine the endocrine physiology/hormone regulation/tissue of origin of rat serum SHBG (part of original tasks 1f and 1g).

D. Methods of Approach

(1) tasks 1(h) and 1(i) --- molecular cloning and cDNA sequencing of horse

SHBG and rat SHBG: Our partial amino acid sequencing of horse SHBG shows it to be only 30 to 40% homologous to human or rabbit SHBG. Although this certainly is a relatively abundant and inexpensive source of SHBG (and CBG), to conduct effective receptor research it is very useful to know the complete amino acid sequence of the ligand. We will do this by cDNA cloning. We will use a horse liver λ ZAP library obtained from Texas A & M. If full length SHBG cDNA cannot be found in this library, fresh horse liver will be used to prepare poly (A⁺) RNA for a new λ ZAP cDNA library by the methods described (69). These expression libraries will be screen for SHBG cDNA with rabbit antibody against human SHBG. It cross reacts with the horse protein although only at relatively low dilutions. The antiserum is available commercially. In addition, we

PI: David A. Sirbasku, Ph.D.

are developing a polymerase chain reaction (PCR) strategy for cloning horse SHBG. From the partial amino acid sequence obtained already, oligonucleotide probes were prepared to amplify specific intervening regions of the mRNA from horse liver poly (A⁺) RNA. This amplified cDNA will be ³²P labeled and used in hybridization screening. This is a newer approach not considered in our original application. Also, rat SHBG in plasma has never been identified. We have now purified a significant supply. If partial amino acid sequencing confirms that rat SHBG is significantly different than SHBGs already reported (70,71), we will conduct the same cDNA cloning study of rat SHBG to determine its complete amino acid sequence. A rat liver library in λ ZAP is available commercially from Stratagene. Since our original application, we have a new fluorescence automated DNA sequencer which will be used in place of the previous manual methods. It determines ≥ 250 bases in a single analysis.

These studies are not necessary with horse CBG because our partial amino acid sequencing shows $\geq 90\%$ homology to the human protein. Also rat CBG has been sequenced previously. The rat CBG isolated by us shows an identical partial sequence to that already published.

(2) work added to TASK 1 by our experimental results--- isolation and cDNA cloning of human Type II SHBG. *We emphasize again that the discovery of a Type II form of SHBG in other species has led to our search of human serum for a similar form. This is an unexpected addition to the project, but certainly one which is central of the original goals of the research program.*

We plan to purify human Type II SHBG from male plasma. Western analysis suggests that males may have relatively higher levels of Type II than females. Indeed, the level of Type II may be lowest in pregnant female serum, which is most often used to isolate SHBG. To purify human Type II SHBG, we will again use the cortisol affinity column method. However, before this is done, Type I SHBG (which has a higher affinity for androgens) will be removed by testosterone affinity chromatography. Additional purification will be achieved by hydrophobic interaction chromatography under various conditions which are expected to separate the Type I and Type II forms.

After completing the purification, we will conduct partial amino acid sequencing by our current methods. The sequences obtained will be used to prepare antibody and oligonucleotide probes for cDNA cloning from human liver libraries. The methods to be employed are described above with horse and rat Type II SHBGs.

(3) tasks 1(d) and 1(e) --- immunological reagents for specifically measuring human Type II SHBG in plasma samples and rat Type II SHBG in rat plasma: The methods for raising rabbit polyclonal antibodies against peptides corresponding to the amino acid sequences of rat SHBG have been described in detail in our original application and will be applied without modification.

For human Type II SHBG, we will use this same methodology with peptides which are clearly different in Types I and II SHBG. This is expected to yield antisera which specifically recognize one or the other of these two forms. If specific identification cannot be achieved by this method, we will use both Type I and Type II human SHBG to raise distinguishing mouse monoclonal antibodies by the methods described in our original application.

PI: David A. Sirbasku, Ph.D.

(4) tasks 1(f) and 1(g) --- endocrine physiology/hormone regulation/tissue of origin of rat serum SHBG: These studies described in our original application will be carried out with only one change. Instead of using mice as our experimental animal, we will use rats. All other methods and procedures will be used without modification. The change in animal model is being made because of our discovery of rat serum Type II SHBG. This change does not in any way alter our original goals.

III. BODY OF THE PROGRESS REPORT

A. Manuscripts in Preparation

The manuscripts listed next are in various stages of preparation and drafting:

- (1) Moreno-Cuevas JE & Sirbasku DA. Steroid hormone-responsive growth of the MTW9/PL2 rat mammary tumor cells in culture medium supplemented with charcoal-dextran extracted serum. (To be submitted to *Molecular and Cellular Endocrinology*)
- (2) Moreno-Cuevas, JE & Sirbasku DA. Estrogen-dependent growth of the MTW9/PL2 rat mammary tumor cells in culture. Evidence for a role of a calcium related inhibitory serum protein. (To be submitted to the *Journal of Steroid Hormone Biochemistry and Molecular Biology Molecular*).
- (3) Tanji, M, Moreno-Cuevas, JE & Sirbasku DA. Purification and characterization of sex hormone-binding globulin (SHBG) and corticosteroid-binding globulin (CBG) from equine serum. (To be submitted to *Endocrinology*).
- (4) Tanji, M, Moreno-Cuevas, JE & Sirbasku DA. Properties of sex hormone-binding globulin (SHBG) and corticosteroid-binding globulin (CBG) purified from adult rat serum by cortisol-agarose and phenyl-Sepharose chromatography. (To be submitted to *Endocrinology*).
- (5) Moreno-Cuevas JE & Sirbasku DA. Identification of sex hormone-binding globulin (SHBG) receptors on MTW9/PL2 rat mammary tumor cells. (To be submitted to *Biochemistry and Biophysics Research Communications*).
- (6) Moreno-Cuevas, JE, Tanji M & Sirbasku DA. Identification of sex hormone-binding globulin (SHBG) as the serum factor which regulates estrogen-responsive MTW9/PL2 rat mammary tumor cell growth in culture. (to be submitted to *Endocrinology*).
- (7) Moreno-Cuevas JE & Sirbasku DA. Sex hormone-binding globulin (SHBG) regulation of estrogen-responsive MTW9/PL2 rat mammary tumor cell growth in serum-free hormonally defined medium. (To be submitted to the *Proceeding of the National Academy of Science, USA*).

PI: David A. Sirbasku, Ph.D.

(8) Moreno-Cuevas JE & Sirbasku DA. Identification of corticosteroid-binding globulin (CBG) receptors on MTW9/PL2 rat mammary tumor cells. (To be submitted to *Molecular and Cellular Endocrinology*).

(9) Moreno-Cuevas JE & Sirbasku DA. Corticosteroid binding-globulin (CBG) regulation of progesterone stimulated MTW9/PL2 rat mammary tumor cell growth in serum-free hormonally defined medium. (To be submitted to *Endocrinology*).

B. Results Obtained During the First 12 Months

We have completed parts of tasks 1(a) through 1(c) and tasks 2(a) and 2(b). These initial studies are summarized next as abstracts of the results to be published in nine manuscripts.

(1) Manuscript 1: The MTW9/PL cell line was established in culture from the carcinogen-induced hormone-responsive MT-W9A rat mammary tumor and shown to form estrogen, androgen and progesterone responsive tumors in Wistar-Furth rats. The MTW9/PL line was used to derive the MTW9/PL2 cells which also exhibit estrogen-responsiveness *in vivo*. In this study, we describe serum supplemented cell culture conditions in which the MTW9/PL2 cell line demonstrates steroid hormone growth responsiveness. The serum used in the cell cultures was steroid hormone-depleted by charcoal-dextran treatment at 34°C. When increasing concentrations of hormone-depleted serum were added to MTW9/PL2 cell cultures, growth was inhibited progressively. Between 30 and 50% (v/v) hormone-depleted serum suppressed growth completely. Addition of 1.0×10^{-13} M to 1.0×10^8 M 17β -estradiol (E_2) reversed the serum dependent inhibition. Under optimal conditions, estrogen induced growth was 2^5 to 2^6 or 32 to 64-fold greater than in control cultures. Even when the cells had been inhibited for six days, E_2 addition restored growth. At 1.0×10^{-11} to 1.0×10^{-7} M, estrone, estriol and the synthetic estrogen diethylstilbestrol promoted optimal growth. Testosterone and dihydrotestosterone were nearly as effective but only at $\geq 10^{-7}$ M. Progesterone was partially effective at $\geq 1.0 \times 10^{-6}$ M. Cortisol was only marginally effective. MTW9/PL2 cells have high affinity receptors for estrogens and progesterone but not androgens. The serum from the adults of several species including rat and human demonstrated activity. Fetal bovine and fetal equine serum showed little activity. The studies described show that the same pattern of hormone-responsiveness seen with MTW9/PL cells in rats is now demonstrable in culture. Furthermore, the MTW9/PL2 cells in culture are the most steroid hormone growth responsive of any of the rat mammary tumor lines yet reported.

Because this cell line so very closely mimics those of human breast, it was used for all of our initial studies of the serum factor identity as SHBG and CBG. Key results were checked with the estrogen and progesterone responsive T47D human breast cancer cell line. This approach saved about six months during the first year. The rat MTW9/PL2 bioassays require 7 days. The T47D assays required 14 days.

PI: David A. Sirbasku, Ph.D.

2. Manuscript 2: The purification of the serum factor mediating the steroid hormone effects on breast cancer cells initially was a very perplexing problem. The factor(s) was completely stable in whole serum for three weeks at 4°C, and yet was remarkably unstable during purification steps lasting only one week. Approximately three months of frustration were invested before we found that simple dialysis caused inactivation. Ultrafiltration proved that the factor was not lost in the dialysate, and therefore, must be a high molecular weight entity. Further exploration revealed that the "factor" was two activities. Their M_r were 200,000 and 55,000. Both were acid and urea labile, and lost when heated at 60°C. Chelex treatment caused inactivation, which suggested metal ion stabilization. A study of metal ions revealed that calcium at 0.1 to 50 mM stabilized the activity. Other metals were not effective. From these constellation of properties, we thought that the activity in serum might be SHBG. Western immunoblotting with anti-human SHBG confirmed that the activity and SHBG coincided in the same fractions from molecular sieve chromatography. However, there were two major dilemmas which contraindicated SHBG. First, we were using horse serum for these studies. Horse serum was not thought to have SHBG. Also, we were assaying with rat mammary cells. Adult rats were not thought to have plasma SHBG. Therefore, we concluded that the calcium stabilized factor(s) was SHBG related immunologically, but was not functionally SHBG.

3. Manuscript 3: Dr. Masao Tanji is an excellent protein chemist trained in this technology in the very prestigious Department of Biochemistry and Biophysics at the University of Tokyo. We are pleased he has come to Houston and joined our group. He analyzed our data and concluded that SHBG in horse serum might be present, but that it had different properties than SHBG from humans or rabbits. His analysis proved correct. In this manuscript in preparation, Dr. Tanji has purified horse serum SHBG and CBG by a simple two step method. This method yielded 5 to 8 mg of SHBG and 26 to 30 mg of CBG per two-liters of horse serum. The molecular size and subunit structure of the two horse proteins was examined by Superdex chromatography, ultracentrifugation and SDS-PAGE \pm reducing agents. The results were consistent with SHBG and CBG. Both proteins were investigated for immunological cross reactivity with anti-human SHBG and anti-human CBG. They cross reacted weakly but specifically by Western analysis. They were examined for steroid hormone binding properties. Horse CBG bound cortisol and progesterone but not dihydrotestosterone or estradiol. Horse SHBG bound dihydrotestosterone and estradiol but not cortisol or progesterone. The partial amino acid sequencing of horse CBG confirmed a high degree of homology with human, rat and rabbit CBGs. The same analysis of horse SHBG showed only 30 to 40% amino acid homologies. We now plan to determine the complete amino acid sequence of horse SHBG. This is an exciting study because the differences and homologies between SHBG from different species can be used to understand such domains as the steroid hormone binding site and the domain(s) which participates in receptor binding. We also have studied the subunit structure of horse SHBG. This was done to gain an additional understanding of how SHBG might interact with cell surface receptors. SHBG has been described as a dimer of identical monomers. Nonetheless, the SDS-PAGE results with SHBG from horse serum showed subunits of M_r 60,000, 52,000 and 25-27,000. Our studies using chemical cross linking followed by ultracentrifugation and SDS-PAGE show that the molecule exists as three dimers representing the possible permutations of the two larger subunits. Furthermore, the 27,000 subunit is actually the two ends of a 52,000 subunit with a

PI: David A. Sirbasku, Ph.D.

proteolytic cleavage in the 164-188 disulfide loop. This cleavage divides the molecular approximately into halves. As yet, we do not know the significance of the subunits nor do we know if the cleavage in this loop has any biological meaning. **However, these studies revealed a potentially major discovery with regard to a new therapy for breast cancer.** The monomers of SHBG are irreversible inhibitors of breast cancer cell growth. Future expansion of this observation is planned and will be the subject of a new NIH grant proposal.

4. Manuscript 4: The issue of the existence of SHBG in adult rat serum has been resolved by our laboratory group. Rats have SHBG, but it is not highly homologous to human or rabbit SHBG. We purchased one liter of a mixture of adult male and female rat serum and used it to conduct the simultaneous purification of CBG and SHBG by the method of Fernlund and Laurell. This yielded 3 mg of CBG and 6 mg of SHBG. The M_r of CBG was 58,000. Only one subunit was found. A partial amino acid sequence of rat CBG was obtained by us and compared to the published structure. There was an exact match in 50 of 50 residues. Next, we studied the rat SHBG. Rat serum SHBG cross reacts with rabbit anti-human SHBG by Western immunoblotting. SDS-PAGE, sedimentation and molecular sieve chromatography confirm a M_r 110,000 dimer. The data support the view that the monomers are very similar. Three subunits of M_r 58,000, 52,000 and 25,000 were identified by SDS-PAGE. Using a filter binding assay, we have found that rat serum SHBG binds dihydrotestosterone and 17β -estradiol, but not cortisol or progesterone. This property defines this molecule as a sex hormone binding globulin. Because rat is well known to express a SHBG-like protein in testis, named rat androgen binding protein (ABP), we compared our SHBG from rat serum to the known amino acid sequence of ABP. By the sequences we have obtained, there is only 50% homology between rat serum SHBG and rat testis ABP. Furthermore, rat serum SHBG is blocked in the amino terminus whereas ABP is not. As stated above, molecular cloning will be used to obtain the complete amino acid sequence of rat serum SHBG.

5. Manuscript 5: We have identified SHBG receptors on MTW9/PL2 rat mammary tumor cells in culture. Radio iodinated SHBG was prepared by a modification of the chloramine T method used by us before. We have established the temperature, time and concentration dependence of ^{125}I -SHBG binding. For these studies specific binding of radio labeled SHBG was measured as total binding minus binding in cultures containing a 300-fold excess of unlabeled SHBG. Binding at 37°C reached a maximum at 2 hours. At 23°C , a maximum was reached at 3 hours. Incubations at 4°C showed a maximum at 6 to 10 hours. At the two higher temperatures, maximum binding was followed by a rapid decline suggesting receptor mediated endocytosis. This was blocked at 4°C . Scatchard analysis indicated a K_d of 1.0 nM for SHBG binding with $\sim 30,000$ sites per cell. All of the studies described were done with ^{125}I -SHBG from which steroid hormone had been removed by charcoal treatment at 34°C . When labeled SHBG was saturated with DHT or estradiol before initiating the binding studies at 4°C , all binding to receptors was blocked. When saturating concentrations of dihydrotestosterone or estradiol were added to the incubation conditions at 37°C , specific binding also was blocked. The data obtained with MTW9/PL2 cells confirm that only steroid hormone depleted SHBG binds to receptors. When sex steroid hormone is bound to free SHBG, the ligand no longer associates with receptor.

PI: David A. Sirbasku, Ph.D.

6. Manuscript 6: In this study, we returned to the characterization of the serum factor(s) regulating estrogen responsive MTW9/PL2 cell growth in culture. As stated above, we knew that the serum factor was calcium related and that horse serum had SHBG. Now, the question became, "is the serum factor regulating estrogen responsiveness actually SHBG". For this study, we first validated a filter membrane method of measuring the SHBG content of serum using ^3H -DHT. This assay was based on a stoichiometry of one mole of ^3H -DHT bound per mole of SHBG dimer. Using this method, we estimated the SHBG concentration of horse serum was 4 to 5 mg/liter. This level compared very favorably to that in human female serum and was higher than in human male. Next, we conducted a series of studies in which we compared the effects of various agents and denaturing conditions on the serum content of SHBG to the effects on estrogen responsive MTW9/PL2 cell growth in culture. First, we measured the SHBG content of serum after dialysis against 0.05 M Tris-HCl, pH 7.4. After 72 hours, all SHBG was lost as measured by the steroid hormone binding assay. This same dialyzed serum also no longer supported estrogenic effects. Based on the fact that calcium and dihydrotestosterone (DHT) bind to SHBG to form a very stable complex, we investigated separately the effects of calcium, DHT and the combination of these on both SHBG stability and the estrogen related activity in serum. To summarize our results, any conditions in which SHBG was inactivated as determined by ^3H -DHT also showed an proportional loss of growth regulating activity. Finally, in this report, we demonstrated that SHBG addition to cultures supplemented with 2.5% horse serum leads to a progressive inhibition of growth which was complete by 2.5 $\mu\text{g/ml}$. Estradiol completely reversed this inhibition thereby restoring growth to maximum rates. The data in this report support the conclusion that SHBG is a major factor in serum which regulates estrogen responsive growth in culture. Nonetheless, the data did not excluded the possibility that something else in serum might be required for SHBG to act as an inhibitor.

7. Manuscript 7: In this report, we extended our study of the regulatory effects of SHBG to completely serum-free conditions. The purpose of this approach was to identify SHBG effects in the complete absence of other potential co-inhibitors in serum. In a previous study, we had reported the development of a serum-free hormonally defined medium for MTW9/PL2 cells. This medium contained F12-DME supplemented with 0.5 mg/ml bovine serum albumin, 10 μM ethanolamine, 10 ng/ml selenous acid, 50 $\mu\text{g/ml}$ apotransferrin, 10 $\mu\text{g/ml}$ diferric transferrin, 10 $\mu\text{g/ml}$ insulin and 1.0 nM T_3 . Growth in this medium was logarithmic for 10 days with a cell population doubling time of ~ 30 hours. The addition of 10 nM 17β -estradiol (E_2) increased the growth rate by 10 to 15%. When SHBG was added, 1.5 to 2.0 $\mu\text{g/ml}$ completely inhibited growth. This response was reversed in part by 1.0×10^{-13} M E_2 . Addition of 1.0×10^{-10} M E_2 promoted optimum growth equal to the estrogen response seen in serum containing cultures. A series of investigations were done to show that horse serum SHBG did not associate with E_2 in culture medium under concentrations reached 500 nM. The experiments presented indeed have confirmed that SHBG is a potent inhibitor of estrogen responsive mammary tumor cell growth and that estrogens reverse the inhibition.

PI: David A. Sirbasku, Ph.D.

8. Manuscript 8: The presence of CBG receptors on MTW9/PL2 rat mammary tumor cells was established by the same methods outlined above for SHBG (see **Manuscript 5**). These studies confirmed approximate 100,000 receptor sites per cell.

9. Manuscript 9: In this report, we have carried out the study of the effects of CBG in serum-free hormonally defined culture. The medium used was the same as in **manuscript 7**. The results of this study were remarkable. Both estrogens and progestins reversed the inhibition caused by CBG. Cortisol had no effect. From the data obtained, we can conclude that both SHBG and CBG are inhibitors of mammary cell growth and that CBG inhibition is different in that progesterone reverses the effect. This will be the first ever report of progesterone regulated breast cancer cell growth in serum-free defined culture.

IV. CONCLUSIONS

	A. MAJOR CONCLUSIONS AND IMPLICATIONS
	(1) SHBG is a physiologically important negative regulator of estrogen responsive breast cancer cell growth. This discovery is important because it is the first to identify a specific serum-borne entity which demonstrates such a function.
	(2) CBG is another major physiologically important negative regulator of progestin and estrogen responsive breast cancer cell growth. This plasma glycoprotein has not previously been recognized as having a function in breast cell growth regulation.
	(3) Our laboratory has identified a new area of breast cancer research. We have established that breast cell growth can be blocked by physiologically available serum-borne inhibitors and that steroid hormone responsive cell types remain quiescent until stimulated by estrogens or progestins.
	(4) Our data raise the possibility that humans may possess a form of SHBG which has not been recognized before. This SHBG (Type II) may be the form responsible for the inhibition of breast tissue growth under normal physiological conditions. We consider it possible that the lack of this form of SHBG in females may be a major predisposing factor to breast cancer.

PI: David A. Sirbasku, Ph.D.

B. FUTURE WORK AND CHANGES TO BETTER ADDRESS THE PROBLEM

There is one problem which is reaching major proportions. Our ability to produce purified SHBG and CBG has become the rate limiting factor in this research. Using the conventional chromatography equipment we have available in the laboratory, these proteins cannot be produced in amounts sufficient to meet our needs.

We have consulted with companies who provide modern automated chromatography systems. Pharmacia has a FPLC (Fast Performance Liquid Chromatography) system which will solve our problem. It is designed to run as an automated preparative system which will produce purified SHBG and CBG at ten day intervals. This system requires about one-quarter of the man-hours we now devote to this effort.

The FPLC will be custom designed for our work. It has computer controlled pumps and valves which move solutions from one column to the next without operator attendance. This permits preparations to proceed overnight and on weekends with only minimal supervision. At present, more than one-half of our laboratory man-hours are devoted to the preparation of these two proteins. This most certainly is not cost effective. In fact, there has been a noticeable slowing of our progress in the last two months because we must stop to make these preparations rather than conducting the next level of experiments.

Mr. Jack Kirten, a Pharmacia bioengineer, is now helping us design an automated FPLC system. His estimate of the costs will be \$27,000 to \$30,000. This amount was not planned into our original application because we had not yet identified the serum factors. Because of this, we could not anticipate the preparative problems we now encounter daily. We are in great need of assistance in this very critical area. As soon as we have a complete/detailed cost estimate from Pharmacia, we will petition the US Army Command for additional funds to purchase this very important equipment.

V. REFERENCES

- (1) Beatson GT (1896) On treatment of inoperable cases of carcinoma of the mamma: suggestions for a new method of treatment with illustrative cases. *Lancet (part 1) July 11*:104-107; *(part 2) July 18*:162-165.
- (2) Jensen EV & DeSombre ER (1972) Mechanism of action of the female sex hormones. *Annu Rev Biochem* 41:203-230.
- (3) O'Malley BW & Means AR (1974) Female steroid hormones and target cell nuclei. *Science* 183:610-620.
- (4) *Laboratory Methods Manual for Techniques in Cellular Signaling and Hormone Action*, Hughes MR, Schrader WT & O'Malley BW (eds), Sixteenth Edition, Houston Biological Associates, Inc.
- (5) Sirbasku DA (1978) Estrogen induction of growth factors specific for hormone-responsive mammary, pituitary, and kidney tumor cells. *Proc Natl Acad Sci USA* 75:3786-3790.
- (6) Ogasawara M, Karey KP, Marquardt H & Sirbasku DA (1989) Identification and purification of truncated insulin-like growth factor I from porcine uterus. Evidence for high biological potency. *Biochemistry* 28:2710-2721.

PI: David A. Sirbasku, Ph.D.

- (7) Karey KP, Marquardt H & Sirbasku DA (1989) Human platelet-derived mitogens. I. Identification of insulinlike growth factors I and II by purification and N^α amino acid sequence analysis. *Blood* 74:1084-1092.
- (8) Riss TL & Sirbasku DA (1989) Identification of a 15,000-molecular weight form of immunoreactive transforming growth factor α in extracts of porcine pituitary. *J Cell Physiol* 138:393-404.
- (9) Riss TL & Sirbasku DA (1989) Characterization of polyclonal antibodies that distinguish acidic and basic fibroblast growth factors by western immunoblotting and enzyme-linked immunosorbent assays. *J Cell Physiol* 138:405-414.
- (10) Karey KP & Sirbasku DA (1988) Differential responsiveness of human breast cancer cell lines MCF-7 and T47D to growth factors and 17 β -estradiol. *Cancer Res* 48:4083-4092.
- (11) Soto AM & Sonnenschein C (1987) Cell proliferation of estrogen-sensitive cells: The case for negative control. *Endocrine Rev* 8:44-52.
- (12) Soto AM, Silva RM & Sonnenschein C (1992) A plasma-borne specific inhibitor of the proliferation of human estrogen sensitive breast tumor cells (estrocologne-I). *J Steroid Biochem Mol Biol* 43:702-712.
- (13) Laursen I, Briand P & Lykkesfeldt AE (1990) Serum albumin as a modulator on growth of the human breast cancer cell line, MCF-7. *Anticancer Res* 10:343-352.
- (14) Dell'Aquila ML, Pigot DA, Bonaquist DL & Gaffney EV (1984) A factor from plasma-derived serum that inhibits the growth of the mammary cell line MCF-7: Characterization and purification. *J Natl Cancer Inst* 72:291-298.
- (15) Furuya Y, Kohno N, Fujiwara Y & Saitoh Y (1989) Mechanism of estrogen action on proliferation of MCF-7 human breast cancer cells in an improved culture medium. *Cancer Res* 49:6670-6674.
- (16) Page MJ, Field JK, Everett NP & Green CD (1983) Serum regulation of the estrogen responsiveness of the human breast cancer cell line MCF-7. *Cancer Res* 43:1244-1250.
- (17) Wiese TE, Kral LG, Dennis KE, Butler WB & Brooks SC (1992) Optimization of estrogen growth response in MCF-7 cells. *In Vitro Cell Dev Biol* 28A:595-602.
- (18) Riss TL & Sirbasku DA (1989) Rat pituitary tumor cells in serum-free culture. II. Serum factor and thyroid hormone requirements for estrogen-responsive growth. *In Vitro Cell Dev Biol* 25:136-142.
- (19) Sorrentino JM, Kirkland WL & Sirbasku DA (1976) Control of cell growth. I. Estrogen-dependent growth in vivo of a rat pituitary tumor cell line. *J Natl Cancer Inst* 56:1149-1154.
- (20) Kirkland WL, Sorrentino JM & Sirbasku DA (1976) Control of cell growth. III. Direct mitogenic effect of thyroid hormones on an estrogen-dependent rat pituitary tumor cell line. *J Natl Cancer Inst* 56:1159-1164.
- (21) Ogasawara M & Sirbasku DA (1988) A new serum-free method of measuring growth factor activities for human breast cancer cells in culture. *In Vitro Cell Dev Biol* 24:911-920.
- (22) Danielpour D, Riss TL, Ogasawara M & Sirbasku DA (1988) Growth of MTW9/PL2 estrogen-responsive rat mammary tumor cells in hormonally defined serum-free media. *In Vitro Cell Dev Biol* 24:42-52.
- (23) Soto AM & Sonnenschein C (1985) The role of estrogens on the proliferation of human breast tumor cells (MCF-7). *J Steroid Biochem* 23:87-94.
- (24) Soto AM, Murai JT, Siiteri PK & Sonnenschein C (1986) Control of cell proliferation: evidence for negative control on estrogen-sensitive T47D human breast cancer cells. *Cancer Res* 46:2271-2275.
- (25) Soto AM, Bass JC & Sonnenschein C (1988) Estrogen-sensitive proliferation pattern of cloned Syrian hamster kidney tumor cells. *Cancer Res* 48:3776-3780.
- (26) Amara JF & Dannies PS (1983) 17 β -estradiol has a biphasic effect on GH cells in culture. *Endocrinology* 112:1141-1143.
- (27) Sato H, Eby JE & Sirbasku DA (1991) Iron is deleterious to hormone-responsive pituitary cell growth in serum-free defined medium. *In Vitro Cell Dev Biol* 27A:599-601.

PI: David A. Sirbasku, Ph.D.

- (28) Sporn MB & Todaro GJ (1980) Autocrine secretion and malignant transformation of cells. *New Engl J Med* 303:878-880.
- (29) Dickson RB, Huff KK, Spenser EM & Lippman ME (1986) Induction of epidermal growth factor related polypeptides by 17 β -estradiol in MCF-7 human breast cancer cells. *Endocrinology* 118:138-142.
- (30) Dickson RB & Lippman ME (1987) Estrogenic regulation of growth and polypeptide growth factor secretion in human breast carcinoma. *Endocrine Rev* 8:29-43.
- (31) Kasid A, Lippman ME, Papageorge AG, Lowry DR & Gelmann EP (1985) Transfection of v-ras^H DNA into MCF-7 cells bypasses their dependence on estrogen for tumorigenicity. *Science* 228:725-728.
- (32) Huff KK, Kaufman D, Gabbay KH, Spencer EM, Lippman ME & Dickson RB (1986) Human breast cancer cells secrete an insulin like growth factor I related polypeptide. *Cancer Res* 46:4613-4619.
- (33) Huff KK, Knabbe C, Lindsey R, Kaufman D, Bronzert D, Lippman ME & Dickson RB (1989) Multihormonal regulation of insulin like growth factor I related protein in MCF-7 breast cancer cells. *Mol Endocrinol* 2:200-208.
- (34) Osborne CK, Coronado EB, Kitten LJ, Arteaga CL, Fuqua SAW, Ramasharma K, Marshal M & Li CH (1989) Insulin like growth factor II (IGF-II): a potential autocrine/paracrine growth factor for human breast cancer acting via the IGF-I receptor. *Mol Endocrinol* 3:1701-1709.
- (35) Kenney NJ, Saeki T, Gottardis M, Kim N, Garcia-Morales P, Martin MB, Normanno N, Ciardiello F, Day A, Cutler ML & Salomon DS (1993) Expression of transforming growth factor α antisense m-RNA inhibits the estrogen-induced production of TGF α and estrogen-induced proliferation of estrogen-responsive human breast cancer cells. *J Cell Physiol* 156:497-514.
- (36) Clarke R, Brunner N, Katz D, Glanz P, Dickson RB, Lippman ME & Kern FG (1989) The effects of constitutive expression of transforming growth factor α on growth of MCF-7 human breast cancer cells *in vitro* and *in vivo*. *Mol Endocrinol* 3:372-380.
- (37) Arteaga CL, Coronado E & Osborne CK (1988) Blockade of the epidermal growth factor receptor inhibits transforming growth factor alpha induced but not estrogen induced growth of hormone dependent human breast cancer. *Mol Endocrinol* 2:1064-1069.
- (38) Arteaga CL & Osborne CK (1989) Growth inhibition of human breast cancer cells in vitro with antibody against the type I somatomedin receptor. *Cancer Res* 49:6237-6241.
- (39) Yee D, Paik S, Lebovic GS, Marcus RR, Favoni RE, Cullen RE, Lippman ME & Rosen N (1989) Analysis of insulin like growth factor I gene expression in malignancy: evidence for a paracrine role in human breast cancer. *Mol Endocrinol* 3:509-517.
- (40) Petra PH (1979) The serum sex steroid binding protein. Purification, characterization and immunological properties of the human and rat proteins. *J Steroid Biochem* 11:245-262.
- (41) Petra PH, Kumar S, Hayes R, Ericksson LH & Titani K (1986) Molecular organization of the sex-steroid binding protein (SBP) of human plasma. *J Steroid Biochem* 24:45-49.
- (42) Rosner W (1990) The functions of corticosteroid-binding globulin and sex hormone-binding globulin: recent advances. *Endocrine Rev* 11:80-91.
- (43) Siiteri PK, Murai JT, Hammond GL, Nisker JA, Raymoure WJ & Kuhn RW (1982) The serum transport of steroid hormones. *Rec Prog Horm Res* 38:457-510.
- (44) Heyns W (1977) The steroid-binding β -globulin of human plasma. *Adv Steroid Biochem Pharmacol* 6:59-79.
- (45) Anderson DC (1974) Sex-hormone-binding globulin. *Clin Endocrinol* 3:69-96.
- (46) Raynaud J-P (1973) Influence of rat estradiol binding plasma protein (EBP) on uterotrophic activity. *Steroids* 21:249-258.

PI: David A. Sirbasku, Ph.D.

- (47) Porto CS, Gunsalus GL, Bardin CW, Phillips DM & Musto NA (1991) Receptor-mediated endocytosis of an extracellular steroid-binding protein (TeBG) in MCF-7 human breast cancer cells. *Endocrinology* 129:436-445.
- (48) Frairia R, Fortunati N, Berta L, Fazzari A, Fissore F & Gaidano G (1991) Sex steroid binding protein (SBP) receptors in estrogen sensitive tissues. *J Steroid Biochem Mol Biol* 40:805-812.
- (49) Fortunati N, Fissore F, Fazzari A, Berta L, Benedusi-Pagliano E & Frairia R (1993) Biological relevance of the interaction between sex steroid binding protein and its specific receptor of MCF-7 cells: effect on the estradiol-induced cell proliferation. *J Steroid Biochem Mol Biol* 45:435-444.
- (50) Strel'chyonok OA, Avvakumov GV & Survilo LI (1984) A recognition system for sex-hormone-binding protein-estradiol complex in human decidual endometrium plasma membranes. *Biochim Biophys Acta* 802:459-466.
- (51) Fortunati N, Fissore F, Fazzari A, Berta L, Giudici M & Frairia R (1991) Sex steroid binding protein interacts with a specific receptor on human pre-menopausal endometrium membrane: modulating effect of estradiol. *Steroids* 56:341-346.
- (52) Porto CS, Abreu LC, Gunsalus GL & Bardin CW (1992) Binding of sex-hormone-binding globulin (SHBG) to testicular membranes and solubilized receptors. *Mol Cell Endocrinol* 89:33-38.
- (53) Hryb DJ, Khan MS & Rosner W (1985) Testosterone-estradiol-binding globulin binds to human prostatic cell membranes. *Biochem Biophys Res Commun* 128:432-440.
- (54) Hryb DJ, Khan MS, Romas NA & Rosner W (1989) Solubilization and partial characterization of the sex hormone-binding globulin receptor from human prostate. *J Biol Chem* 264:5378-5383.
- (55) Nakhla AF, Khan MS & Rosner W (1990) Biologically active steroids activate receptor-bound human sex hormone-binding globulin to cause LNCaP cells to accumulate adenosine 3',5'-monophosphate. *J Clin Endocrinol Metab* 71:398-404.
- (56) Khan MS, Hryb DJ, Hashim GA, Romas NA & Rosner W (1990) Delineation and synthesis of the membrane receptor-binding domain of sex hormone-binding globulin. *J Biol Chem* 265:18362-18365.
- (57) Hryb DJ, Khan MS, Romas NA & Rosner W (1990) The control of the interaction of sex hormone-binding globulin with its receptor by steroid hormones. *J Biol Chem* 265:6048-6054.
- (58) Rosner W, Hryb DJ, Khan MS, Singer CJ & Nakhla AM (1988) Are corticosteroid-binding globulin and sex hormone-binding globulin hormones? *Ann NY Acad Sci* 538:137-145.
- (59) Nakhla AM, Khan MS & Rosner W (1988) Induction of adenylate cyclase in a mammary carcinoma cell line by human corticosteroid-binding globulin. *Biochem Biophys Res Commun* 153:1012-1018.
- (60) Rosner W, Hryb DJ, Khan S, Nakhla AM & Romas NA (1991) Sex hormone-binding globulin: anatomy and physiology of a new regulatory system. *J Steroid Biochem Mol Biol* 40:813-820.
- (61) Kuhn RW (1988) Corticosteroid-binding globulin interactions with target cells and plasma membranes. *Ann NY Acad Sci* 538:146-158.
- (62) Castilla LH, Couch FJ, Erdos MR, Hoskins KF, Calzone K, Garber JE, Boyd J, Lubin MB, Deshano ML, Brody LC, Collins FS & Weber BL (1994) Mutations in the BRCA1 gene in families with early-onset breast and ovarian cancer. *Nature Genetics* 8:387-391.
- (63) Shattuck-Eidens D, McClure M, Simard J, Labrie F, Narod S, Couch F, Hoskins K, Weber B, Castilla L, Erdos M, Brody L, Friedman L, Ostermeyer E, Szabo C, King MC, Jhanwar S, Offit K, Norton L, Gilewski T, Lubin M, Osborne M, Black D, Boyd M, Steel M, Ingles S, Haile R, Lindblom A, Olsson H, Borg A, Bishop DT, Solomon E, Radice P, Spatti G, Gayther S, Ponder B, Warren W, Stratton M, Liu Q, Fujimura F, Lewis C, Skolnick MH & Goldgar DE (1995) A collaborative survey of 80 mutations in the BRCA1 breast and ovarian cancer susceptibility gene. *JAMA* 273:535-541.

PI: David A. Sirbasku, Ph.D.

- (64) Smith SA, Easton DF, Evans DGR & Ponder BAJ (1992) Allele losses in the region 17q12-21 in familial breast and ovarian cancer involve the wild type chromosome. *Nature Genetics* 2:128-131.
- (65) Elledge RM, Fuqua SAW, Clark JM, Pujol P & Allred DC (1993) The role and prognostic significance of p53 gene alterations in breast cancer. *Breast Cancer Res Treat* 27:95-102.
- (66) Elledge RM & Allred DC (1994) The p53 tumor suppressor gene in breast cancer. *Breast Cancer Res Treat* 32:39-47.
- (67) Jolly KW, Malkin D, Douglas EC, Brown TF, Sinclair AE & Look AT (1994) Splice-site mutation of the p53 gene in a family with hereditary breast-ovarian cancer. *Oncogene* 9:97-102.
- (68) Malkin D, Li FP, Stong LC, Fraumeni JF, Nelson CE, Kim DH, Kassel J, Gryka MA, Bischoff FZ, Tainsky MA & Friend SH (1990) Germline p53 mutations in a familial syndrome of breast cancer, sarcomas and other neoplasms. *Science* 250:1233-1238.
- (69) Davis LG, Dibner MD & Battey JF (1986) Basic Methods in Molecular Biology, Elsevier, New York.
- pp 72-74 -- End labeling of synthetic probes.
 - pp 75-78 -- Hybridization with synthetic ³²P end-labeled probe.
 - pp 130-135 -- Guanidine isothiocyanate preparation of total RNA.
 - pp 143-146 -- Formaldehyde gel for electrophoretic separation of RNA and Northern blot
 - pp 199-207 -- Generation of cDNA insert from eukaryotic mRNA.
 - pp 211-215 -- Plating and screening of λ gt10 and λ gt11 packaged inserts.
 - pp 80-83 -- Nick translation.
 - pp 276-284 -- S1 nuclease protection assay.
- (70) Petra PH, Que BG, Namkung PC, Ross JBA, Charbonneau H, Walsh KA, Griffin PR, Shabanowitz J & Hunt DF (1988) Affinity labeling, molecular cloning, comparative amino acid sequence analyses of sex steroid-binding protein of plasma. *NY Acad Sci* 538:10-24.
- (71) Walsh KA, Titani K, Kumar S, Hayes R & Petra PH (1986) Amino acid sequence of the sex steroid binding protein of human plasma. *Biochemistry* 25:7584-7590.